



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Statistical reports of States and cities of the United States—Yearly and monthly.*

CONNECTICUT—*Bridgeport*.—Month of February, 1901. Estimated population, 71,000. Total number of deaths, 112, including scarlet fever, 4, and 16 from tuberculosis.

INDIANA.—The total number of deaths reported for the month of February, 1901, was 3,350, and the death rate is 17.3. For the corresponding month of last year the deaths numbered 2,732, a rate of 13.4. The deaths during the month under 1 year of age were 566; 1 to 5 inclusive, 214, and 65 years and over, 951. The deaths from all forms of pulmonary tuberculosis were 411 as against 395 for February, 1900. The rate from these figures is 193.2 per 100,000. Typhoid fever caused 45 deaths; diphtheria, 55; scarlet fever, 23; measles, 19; whooping cough, 27; pneumonia, 608; diarrheal diseases, 8; cerebro-spinal meningitis, 43; influenza, 324; cancer, 86; violence, 96, and smallpox, 2. Pulmonary consumption, pneumonia, and influenza account for 39 per cent of all deaths. A curiosity in the death table is the fact that the deaths from whooping cough were greater than from scarlet fever, the figures being, respectively, 27 and 23. Cancer seems to be slowly on the increase.

MASSACHUSETTS—*Fitchburg*.—Month of January, 1901. Census population, 31,531. Total number of deaths, 46, including diphtheria, 2, and 5 from phthisis pulmonalis.

Month of February, 1901. Total number of deaths, 42, including diphtheria, 2; la grippe, 3, and 1 from phthisis pulmonalis.

*Worcester*.—Month of February, 1901. Census population, 118,421. Total number of deaths, 175, including diphtheria, 2; enteric fever, 3; scarlet fever, 5; whooping cough, 1, and 26 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended March 9, 1901, from 83 observers indicate that inflammation of kidneys, scarlet fever, erysipelas, inflammation of bowels, diphtheria, smallpox, dysentery, and measles were more prevalent, and whooping cough was less prevalent than in the preceding week.

Cerebro-spinal meningitis was reported present at 2, whooping cough at 12, measles at 29, diphtheria at 33, enteric fever at 45, scarlet fever at 110, phthisis pulmonalis at 176, and smallpox at 72 places.

*Grand Rapids*.—Month of February, 1901. Estimated population, 90,000. Total number of deaths, 106, including diphtheria, 2; enteric fever, 5; scarlet fever, 3; la grippe, 7, and 7 from tuberculosis.

The Monthly Bulletin of Vital Statistics for February, 1901, says:

There were 3,145 deaths returned as having occurred in Michigan during February. This number is 28 less than the number returned for January, but owing to the shortness of the month, the death rate per thousand population increased from 15.7 to 17.1. There were 472 more deaths in February, 1901, than in February, 1900.

By ages, there were 536 deaths of infants under 1 year of age; 185 deaths of children aged 1 to 4 years, and 1,029 deaths of persons aged 65 years and over.

Important causes of death were as follows: Pulmonary tuberculosis, 165; other forms of tuberculosis, 19; typhoid fever, 43; diphtheria and croup, 47; scarlet fever, 34; measles, 5; whooping cough, 13; pneumonia, 474; influenza, 378; cancer, 121; accidents and violence, 100.

There was a marked decline in tuberculous diseases, typhoid fever, diphtheria, and croup, and a very considerable increase in the number of deaths reported from influenza. Pneumonia also increased, but not to as large an extent.

There was one death reported from smallpox in the city of Saginaw. The number of deaths in Indiana for the month of February was 3,350, corresponding to a death rate of 17.3 per 1,000 population.

MINNESOTA—*Minneapolis*.—Month of January, 1901. Census population, 202,718. Total number of deaths, 250, including diphtheria, 24; enteric fever, 5; scarlet fever, 3; whooping cough, 1, and 31 from tuberculosis.

Month of February, 1901. Total number of deaths, 192, including diphtheria, 7; enteric fever, 4; scarlet fever, 1; whooping cough, 2, and 22 from tuberculosis.

*St. Paul*.—Month of February, 1901. Estimated population, 165,000. Total number of deaths, 122, including diphtheria, 1; enteric fever, 2; measles, 1; scarlet fever, 2, and 13 from tuberculosis.

OHIO.—Reports to the State board of health for the eight weeks ended February 23, 1901, from 69 localities having an aggregate estimated population of 1,363,503, show 100 deaths from diphtheria, 98 from enteric fever, 5 from measles, 21 from scarlet fever, and 4 from whooping cough.

*Cleveland*.—Month of February, 1901. Estimated population, 395,000. Total number of deaths, 461, including diphtheria, 18; enteric fever, 17; scarlet fever, 4, and 28 from tuberculosis.

PENNSYLVANIA—*Oil City*.—Month of January, 1901. Estimated population, 13,364. Total number of deaths, 17, including diphtheria, 1; enteric fever, 1, and 2 from tuberculosis.

Month of February, 1900. Total number of deaths, 23, including enteric fever, 2, and 1 from tuberculosis.

TENNESSEE—*Nashville*.—Month of February, 1901. Census population, 80,865—white, 50,627; colored, 30,238. Total number of deaths, 155, including enteric fever, 3; whooping cough, 2, and 27 from tuberculosis.

UTAH—*Ogden*.—Month of February, 1901. Census population, 16,313. Total number of deaths, 9. No deaths from contagious diseases.

WASHINGTON—*Seattle*.—Month of February, 1901. Estimated population, 90,000. Total number of deaths, 73, including diphtheria, 1; enteric fever, 1; scarlet fever, 1, and 7 from tuberculosis.

*Report of immigration at Boston for the week ended March 16, 1901.*

OFFICE OF U. S. COMMISSIONER OF IMMIGRATION,  
*Port of Boston, March 18, 1901.*

*Number of alien immigrants who arrived at this port during the week ended March 16, 1901; also names of vessels and ports from which they came.*

Date.	Vessel.	Where from.	No. of immigrants.
Mar. 10	Steamship Prince Arthur .....	Yarmouth, Nova Scotia.....	104
Do....	Steamship Boston.....	.....do.....	100
Do....	Steamship Admiral Dewey .....	Port Morant, Jamaica .....	1
Mar. 12	Steamship Ultonia .....	Liverpool, England .....	157
Mar. 13	Steamship English King .....	Antwerp, Belgium .....	1
Do....	Steamship Memnon .....	Avonmouth, England.....	1
Mar. 14	Steamship Pro Patria.....	St. Pierre, Miquelon.....	2
Do....	Steamship Prince George.....	Yarmouth, Nova Scotia.....	48
Do....	Steamship Boston.....	.....do.....	91
Do....	Steamship Halifax.....	Halifax, Nova Scotia .....	32
Mar. 16	Steamship Bostonian.....	London, England .....	3
Do....	Steamship Bohemian .....	Liverpool, England .....	1
	Total.....	.....	541

GEORGE B. BILLINGS, *Commissioner.*